No.



9000028

THE UNIVERD STAYIES OF ANTERIOR

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Minnesota Agricultural Experiment Station

Colherens, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE; IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANTS VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S), AND THE SUCCESSORS, HEIRS OF ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF "Cighteen" TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITOR) AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT RIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT.

THE UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS BY THE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

SOYBEAN

'Kato'

In Eastmany Expersor, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington, D.C. this 31st day of December in the year of our Lord one thousand nine hundred and ninety-one.

Kennth Hevan Commissioner Plant Variety Grotection Office Spainthwest Marketing Conjection

Shoped MAdig In Secretary of Agriculture

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Office, DIRM, Room 404-W, Washington, D.C. 20250; and to the Office of Management and Budget, Paperwork Reduction Project (DMB #081-0055), Washington, 20250.

	Table of the state	·		0001 0000; Expires 110 1/01
APPLICATION FOR PLANT VARIETY	PROTECTION	CERTIFICATE	de ce In	stermine if a plant variety protection rtificate is to be issued (7 U.S.C. 2421). formation is held confidential until
	3.007	2. TEMPORARY DESIGNATION	NOR 3	VARIETY NAME
	ion	EXPERIMENTAL NO. M81-382		Kato
4. ADDRESS (street and no. or R.F.D. no., city, state, and ZIP)		5. PHONE (Include area code)	FOR OFFICIAL USE ONLY
University of Minnesota			PV	
220 Coffey Hall				9000028
		612/625-4221		7000020
St. Paul, MN 55108				4
6. GENUS AND SPECIES NAME 7	FAMILY NAME (Botanic	at)	;	16 1489
4	•	a.,		·
		ATE OF DETERMINATION		
·	1	_		3 270 0,
		•		Date
	CHON (Corporation, parti	ersnip, association, etc.)		
· · · · · · · · · · · · · · · · · · ·	[
TIL IF INCORPORATED, GIVE STATE OF INCORPORATION	12. DA	E OF INCORPORATION		
13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY, TO SER	Innesota Agricul tural Experiment Station M81-382 Kato DOPRESS datived and no. or R.P.D. no. eths. states. and 2019 Inversity of Minnesota 20 Coffey Hall 420 Eckles Avenue t. Paul, NN 55108 ENUS AND SPECIES NAME I. Paul, NN 59, 1988 I. DATE OF DETERMINATION I. DATE OF DETERMINATION I. DATE OF DETERMINATION II. DATE OF DETERMINATION II. DATE OF DETERMINATION II. DATE OF DETERMINATION III. DATE OF DECOMPORATION III. DATE OF DECOMPO			
University of Minnesota, 1991 Buford Ci 411 Borlaug Hall St. Paul, MN 55108	ircle	PHONE (Include a	rea code):	
r us	NOTITIONS OF TEVERS			
d. Exhibit D, Additional Description of Variety.				
e. X Exhibit E, Statement of the Basis of Applicant's Ownership.				
f X Seed Sample (2,500 viable untreated seeds). Date Seed Sam	nple mailed to Plant Va	ariety Protection Office		
and the same of th				
16. DOES THE APPLICANT(S) SPECIFY THAT THIS VARIETY BE LIMITED AS TO NUMBER OF GENERATIONS?	i 17. IF "YES" TO	ITEM 16, WHICH CLASSES OF	PRODUCTIO	N BEYOND BREEDER SEED?
X YES NO	X FOU	IDATION	REGISTEREE	CERTIFIED
18. DID THE APPLICANT(S) PREVIOUSLY FILE FOR PROTECTION OF THE VARIET	Y IN THE U.S.?	·		
X NO	-			
19. HAS THE VARIETY BEEN RELEASED, USED, OFFERED FOR SALE, OR MARKI	ETED IN THE U.S. OR O	THER COUNTRIES?	· · 	
	•	•		
X NO				
20. The applicant(s) declare(s) that a viable sample of basic seeds or request in accordance with such regulations as may be applicab	of this variety will l	pe furnished with the app	lication ar	nd will be replenished upon
uniform, and stable as required in section 41, and is entitled to	protection under th	e provisions of section 42	of the Plan	hat the variety is distinct, t Variety Protection Act.
Applicant(s) is (are) informed that false representation herein c	can jeopardize prote	ction and result in penalt	es.	
SIGNATURE OF APPLICANT (Owner(s))	CAPACITY OR TI	TLE	T	DATE
Jour hampson	Assista	nt Director		November 10, 1989
SIGNATURE OF APPLICANT (Owner(s))	CAPACITY OR TI	TLE		DATE
L'Engene Allen	Acting 1	Director		November 10, 1989
FORM CSSD-470 (5.89) Edition of FORM LS-470, 3-86, is obsolete.		The second second		

Exhibit A

Origin and Breeding History of Kato Soybean

'Kato' traces to the F_5 progeny of an F_4 plant pulled from a population that had been advanced by a modified single seed descent procedure from the cross M70-127 x Century. M70-127 is a line derived from the cross Evans x M63-217Y; M63-217Y is a sib of the cultivar Hodgson having yellow hilum. Bulked seed of the F5 row was designated M81-382 and was used for yield testing in the F_6 (1982). Subsequent tests of strain M81-382 were conducted in the F_7 (1983) and F_8 (1984). In the F_8 generation, 50 typical plants were selected to initiate purification for observable traits including reaction to race 1 of Phytophthora root rot. In the F_Q (1985), M81-382 was entered in the Maturity Group I Preliminary Regional Soybean Test. In 1985, thirty-nine rows were grown for purification purposes. Twenty-one rows appeared uniform for plant and seed characteristics including resistance to race 1 of Phytophthora root rot, therefore, seed of these rows were bulked to provide the breeders' seed. In the F_{10} (1986), F_{11} (1987) and F_{12} (1988), M81-382 was tested in the Uniform Regional Soybean Test Maturity Group I. In the F_{10} (1986), a small increase of breeders' seed was made. In the F_{11} (1987), foundation seed was produced by the Minnesota Foundation Seeds Organization. The foundation seed produced was shared with other states for increase. In the F_{12} (1988), seed was increased and M81-382 was approved for release as Kato. On February 15, 1989, seed of Kato was released to registered and/or certified growers in Minnesota and South Dakota. No off type variants were noted in the seed multiplication process of Kato over three generations, thus the variety breeds true and meets certification standards.

Exhibit B

25612 625 1268

Novelty Statement

'Kato' is most similar to 'Sibley'. Kato has purple flowers versus white flowers for Sibley. Kato has tawny pubescence while Sibley has gray pubescence. The hilum color for Kato is black while the hilum color for Sibley is yellow. Kato matures approximately the same time as Sibley, has about the same yield potential and is about two inches taller. Kato has larger seeds than Sibley. Kato has a better plant lodging score but a poorer seed quality score than Sibley. The protein content of Kato is considerably higher than Sibley but the oil content of Kato is lower than the oil content of Sibley. Both varieties have the Ros1 gene for Phytophthora root rot resistance.

Data comparing Kato and Sibley is taken from Uniform Test I Northern States 1986-88 (a total of 43 tests for most traits).

					Seed			
Variety	Date mature	Yield bu/ac	Height inches	Lodging score	quality score	Seed size	Oil %	Protein %
Kato	9/14	42.2	36	1.5	2.5	19.5	19.7	42.1
Sibley	9/14	42.0	34	2.1	2.0	17.3	21.4	39.7

EXHIBIT C (Soybean)

U.S. DEPARTMENT OF AGRICULTURE
AGRICULTURAL MARKETING SERVICE
LIVESTOCK, MEAT, GRAIN & SEED DIVISION
PLANT VARIETY PROTECTION OFFICE
BELTSVILLE, MARYLAND 20705

OBJECTIVE DESCRIPTION OF VARIETY SOYBEAN (Glycine max L.)

SUYBE	AN (GIYCINE	max L./		
NAME OF APPLICANT(S)	TEMPORARY	DESIGNATION VARIE	TY NAME	
linnesota Agricultural Experiment Station	M82-382	K	ato	
ADDRESS (Street and No., or R.F.D. No., City, State, and Zip Co University of Minnesota 220 Coffey Hall, 1420 Eckles Avenue St. Paul, MN 55108	de)	PVPO	NUMBER	00028
Choose the appropriate response which characterizes the vain your answer is fewer than the number of boxes provided				
1. SEED SHAPE: 2 1 = Spherical (L/W, L/T, and T/W ratios = < 1.2) 3 = Elongate (L/T ratio > 1.2; T/W = < 1.2)	1 1 2 = \$ p	herical Flattened (L/W ra ongate Flattened (L/T ra		
2. SEED COAT COLOR: (Mature Seed)				
1 = Yellow 2 = Green 3 = Brown	4 = Black	5 = Other (Specif	γ)	
3. SEED COAT LUSTER: (Mature Hand Shelled Seed)				
1 = Dull ('Corsoy 79'; 'Braxton') 2 = Shiny ('Neb	soy'; 'Gasoy 17')	<u></u>		
4. SEED SIZE: (Mature Seed)				·
1 9 Grams per 100 seeds	·			
5. HILUM COLOR: (Mature Seed)				
6 1 = Buff 2 = Yellow 3 = Brown	4 = Gray	5 = Imperfect Black	6 = Black	7 = Other (Specify)
6. COTYLEDON COLOR: (Mature Seed)				
1 = Yellow 2 = Green				
7. SEED PROTEIN PEROXIDASE ACTIVITY:				
2 1 = Low 2 = High				
8. SEED PROTEIN ELECTROPHORETIC BAND:	,			
2 = Type A (SP1 ^a) 2 = Type B (SP1 ^b)				·
9. HYPOCOTYL COLOR:				
1 = Green only ('Evans'; 'Davis') 2 = Green w 3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71' 4 = Dark Purple extending to unifoliate leaves ('Hodgson')	")	elow cotyledons ('Woodv	vorth'; 'Tracy')	
10. LEAFLET SHAPE:				
3 1 = Lanceolate 2 = Oval 3 = Ovate	e 4 = Ot	her (Specify)		

11. LEA	FLET SIZE:			
2	1 = Small ('Amsoy 71'; 'A5312') 3 = Large ('Crawford'; 'Tracy')	2 = Medium ('Corsoy 79'; 'Gasoy 17')		
40 154	F COLOR:			
2. LEA	1 = Light Green ('Weber'; 'York') 3 = Dark Green ('Gnome'; 'Tracy')	2 = Medium Green ('Corsoy 79'; 'Braxto	n r')	:
13. FLO	WER COLOR:		470.0	
2	1 = White 2 = Purple	3 = White with purple throat		
14. POD	COLOR:			
2	1 = Tan 2 = Brown	3 = Black	.1.	
15. PLAI	NT PUBESCENCE COLOR:	*		
2	1 = Gray 2 = Brown (Tawny)			
16. PLAN	IT TYPES:			
2	1 = Slender ('Essex'; 'Amsoy 71') 3 = Bushy ('Gnome'; 'Govan')	2 = Intermediate ('Amcor'; 'Braxton')		
	**			
17. PLAN	IT HABIT:			
3 1	1 = Determinate ('Gnome'; 'Braxton')	2 = Semi-Determinate ('Will')		•
\$14	; 3 = Indeterminate ('Nebsoy'; 'Improved Pe	lican')		•
\$N	JRITY GROUP:	lican')		
\$N	0	4 = I 5 = II 6 = III	7 = IV 8 = V	
18. MATU 0 4	URITY GROUP: 1 = 000	4 = I 5 = II 6 = III I 12 = IX 13 = X	7 = IV 8 = V	
18. MATU 0 4 19. DISEA	### COUP: 1 = 000	4 = I 5 = II 6 = III I 12 = IX 13 = X	7 = IV 8 = V	
18. MATU 19. DISEA BAC	### COUP: 1 = 000	4 = I 5 = II 6 = III I 12 = IX 13 = X Susceptible; 2 = Resistant)	7 = IV 8 = V	
18. MATU 0 4 19. DISEA	### COUP: 1 = 000	4 = I 5 = II 6 = III I 12 = IX 13 = X Susceptible; 2 = Resistant)	7 = IV 8 = V	
18. MATU 19. DISEA BAC	### COUP: 1 = 000	4 = I 5 = II 6 = III I 12 = IX 13 = X Susceptible; 2 = Resistant)	7 = IV 8 = V	
18. MATU 19. DISEA BAC	JRITY GROUP: 1 = 000	4 = I 5 = II 6 = III I 12 = IX 13 = X Susceptible; 2 = Resistant)	7 = IV 8 = V	
18. MATU 0 4 19. DISEA BAC 0 0	JRITY GROUP: 1 = 000	4 = I 5 = II 6 = III I 12 = IX 13 = X Susceptible; 2 = Resistant)	7 = IV 8 = V	
18. MATU 0 4 19. DISEA BAC 0 0	JRITY GROUP: 1 = 000	4 = I 5 = II 6 = III I 12 = IX 13 = X Susceptible; 2 = Resistant)	7 = IV 8 = V	
18. MATU 19. DISEA BAC 0 0 FUNG	JRITY GROUP: 1 = 000	4 = I 5 = II 6 = III I 12 = IX 13 = X Susceptible; 2 = Resistant)	7 = IV 8 = V	
18. MATU 19. DISEA BAC 0 0 FUNG	JRITY GROUP: 1 = 000	4 = I 5 = II 6 = III I 12 = IX 13 = X Susceptible; 2 = Resistant)	7 = IV 8 = V Other (Specify)	
18. MATU 19. DISEA BAC 0 0 FUNG	JRITY GROUP: 1 = 000	4 = I 5 = II 6 = III I 12 = IX 13 = X Susceptible; 2 = Resistant) ar. sojensis)		
18. MATU 19. DISEA BAC 0 0 FUNG 0	JRITY GROUP: 1 = 000	4 = I		
18. MATU 19. DISEA BAC 0 0 FUNG 0	JRITY GROUP: 1 = 000	4 = I		
18. MATU 19. DISEA BAC 0 0 FUNG 0	JRITY GROUP: 1 = 000	4 = I		

19,	DISEA	SE REACTION	I: (Enter 0 = Not Test	ed; 1 = Susceptible; 2 =	Resistant) (C	ontinued)		
•	FUN	GAL DISEASE	S: (Continued)					
	0	Pod and Sten	n Blight <i>(Diaporthe pha</i>	seolorum var; sojae)		•		
	0	Purple Seed S	Stain <i>(Cercospora kikud</i>	hii)				
	0	Rhizoctonia	Root Rot (Rhizoctonia	solani)				
		Phytophthora	a Rot (Phytophthora m	egasperma var. sojae)				
	2	Race 1	0 Race 2	Race 3	Race 4	0 Race 5	0 Race 6	0 Race 7
	0	Race 8	0 Race 9	Other (Specify)				
	VIRA	AL DISEASES:						
	0	Bud Blight (T	obacco Ringspot Virus)				•
	0	Yellow Mosai	c (Bean Yellow Mosaic	Virus)				•
	0	Cowpea Mosa	nic (Cowpea Chlorotic '	/irus)				
	0	Pod Mottle (E	Bean Pod Mottle Virus)					
	0	Seed Mottle (Soybean Mosaic Virus)					
	NEM	ATODE DISEA	ASES:			•		
		Soybean Cyst	Nematode (Heteroder	a glycines)	1,1 .			
	1	Race 1	0 Race 2	Race 3	Race 4	Other (S)	pecify)	
	0	Lance Nemate	ode (Hoplolaimus Colo	mbus)	•			
	0	Southern Roo	ot Knot Nematode (Me	oidogyne incognita)	# 2"			
		Northern Roo	ot Knot Nematode (Me	oidogyne Hapla)				
		Peanut Root I	Knot Nematode <i>(Meloi</i>	dogyne arenaria)				
		Reniform Nen	natode (<i>Rotylenchulus</i>	reniformis)	•			
-	\sqcap	OTHER DISE	ASE NOT ON FORM	(Specify):	·			
			 			····		
0. 1		LOGICAL RES	SPONSES: (Enter 0 =	Not Tested; 1 = Suscept	tible; 2 = Res	stant)		·
	2	Iron Chlorosis	on Calcareous Soil		٠			
		Other (Specify	v)		•			
1. 1	NSECT	REACTION:	(Enter 0 = Not Tested;	1 = Susceptible; 2 = Re	sistant)			
	0	Mexican Bean	Beetle (Epilachna vari	restis)				
	0	Potato Leaf H	opper <i>(Empoasca faba</i> e	<i>y</i>	ē			
	0	Other (Specify	<i>y</i>		· · · · · · · · · · · · · · · · · · ·			
2 1	NDICAT	TE WHICH VA	RIETY MOST CLOSE	LY RESEMBLES THA	T SUBMITTE	D.		· · · · · · · · · · · · · · · · · · ·
	CHAR	ACTER	NAME O	VARIETY	CHAI	RACTER	NAME	OF VARIETY
Р	lant Sha	pe	Hodgson 78		Seed Co	at Luster	Sibley	
L	eaf Shap	oe .	Hodgson 78		Seed Siz	e	Sibley	
L	eaf Colo	or	Hodgson 78		Seed Sh	эре	Sibley	<u> </u>
L	eaf Size		Hodgson 78	· · · · · · · · · · · · · · · · · · ·	Seedling	Pigmentation	Hodgson 78	·
			'					

23. GIVE DATA FOR SUBMITTED AND SIMILAR STANDARD VARIETY: Paired Comparison Data

VARIETY	NO. OF DAYS MATURITY	PLANT LODGING SCORE	CM PLANT HEIGHT	LEAFLET SIZE		SEED CONTENT		SEED SIZE G/100	NO. SEEDS/
				CM Width	CM Length	% Protein	% Oil	SEEDS	POD
Kato Submitted	120	1.5	91	9.2	11.7	42.1	19.7	19.5	2.4
Sibley Name of Similar Variety	120	2.1	86	9.8	11.9	39.7	21.4	17.3	2.4

PUBLICATIONS USEFUL AS REFERENCE AIDS FOR COMPLETING THIS FORM:

- 1. Caldwell, B.E., ed. 1973. Soybeans: Improvement, Production, and Uses. Amer. Soc. Agron. Monograph No. 16.
- 2. Buttery, B.R. and R.I. Buzzell. 1968. Peroxidase activity in seeds of soybean varieties. Crop Sci., 8: 722-725.
- 3. Hymowitz, T. 1973. Electrophoretic analysis of SBTI-A2 in the USDA soybean germplasm collection. Crop Sci., 13: 420-421.
- 4. Payne, R.C. and L.F. Morris. 1976. Differentiation of soybean cultivars by seedling pigmentation patterns. J. Seed Technol. 1: 1-19.



Exhibit E

Statement of the Basis of Ownership

The Minnesota Agricultural Experiment Station is the owner of Kato. The Minnesota Agricultural Experiment Station is the employer of the breeders who developed Kato.